

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>A61K 47/40, 47/48, 31/565</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 97/17992</b> <b>(43) International Publication Date:</b> 22 May 1997 (22.05.97)
<b>(21) International Application Number:</b> PCT/US96/18189 <b>(22) International Filing Date:</b> 12 November 1996 (12.11.96)  <b>(30) Priority Data:</b> 06/006,557 13 November 1995 (13.11.95) US Not furnished 7 November 1996 (07.11.96) US  <b>(71) Applicant:</b> SUPERGEN, INC. [US/US]; 6450 Hollis Street, Emeryville, CA 94608 (US).  <b>(72) Inventors:</b> RUBINFELD, Joseph; 5304 Blackhawk Drive, Danville, CA 94506 (US). VIDA, Julius, A.; 27 Sachem Road, Greenwich, CT 06830 (US). BRADLOW, H., Leon; 86-25 Palo Alto Street, Holliswood, NY 11423-1203 (US). FINEMAN, Elliott, L.; 68 Stratford Road, Kensington, CA 94707 (US).  <b>(74) Agent:</b> FINEMAN, Elliott, L.; 68 Stratford Road, Kensington, CA 94707 (US).		<b>(81) Designated States:</b> AU, CA, CN, HU, IL, JP, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).  <b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
<b>(54) Title:</b> IMPROVED FORMULATION FOR ADMINISTRATION OF STEROID COMPOUNDS  <b>(57) Abstract</b>  Improved cyclodextrin formulations for the administration of 5 $\beta$ steroids including $\alpha$ etiocholanolone, $\beta$ etiocholanolone, and etiocholandione are provided. Also provided are cyclodextrin formulations including at least one of the 5 $\beta$ steroids and DHEA. The formulations are suitable for parenteral administration, particularly intravenous administration and are also suitable for treating a condition which responds to treatment with the 5 $\beta$ steroids alone or together with DHEA, particularly where it is desired to decrease the amount of DHEA administered, including such conditions as obesity, diabetes syndrome, diabetes-associated hypercorticism or combinations thereof, anemic disorders including but not limited to aplastic anemia and anemia associated with renal failure or chemotherapy-induced or radiation-induced anemia or neutropenia, and autoimmune or inflammatory disorders such as lupus erythematosus.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

## Improved Formulation for Administration of Steroid Compounds

### Field of the Invention

5

The present invention relates to improved formulations for the administration of certain steroid and steroid related compounds, which renders them more effective and which also facilitates parenteral administration of these compounds.

### 10 Background of the Invention

Steroid compounds are powerful drug substances generally well known in biology, medicine and pharmaceutical sciences. These compounds effect a number of important physiological functions in mammals and in human beings in particular.

15

Many of the beneficial steroids that are administered therapeutically to patients are insoluble in water. As a result, few of the active steroids can be administered in aqueous solution parenterally. As a result of the insolubility of steroids in aqueous media many of the active steroids are administered in organic solvent vehicles particularly when administered topically and by injection. If injected, steroids are usually administered using a needle into the muscle or intramuscularly (im) or into the layer below the skin or subcutaneously. Steroids injected by both of these routes are absorbed gradually into the blood of the patient. The resulting peak concentration of the drug in the blood is obtained only slowly and the therapeutic effect of the drug may be achieved only gradually and after some delay.

25

Because of these problems in parenteral administration of steroids, these compounds are usually administered orally especially when chronic administration is required to exert a therapeutic effect. Absorption of the compounds by the oral route however, is limited. In most cases, if taken orally, less than 15 percent of the drug reaches the blood stream. One reason for the low percentage of steroid reaching the blood stream is that much of the drug that is absorbed across the intestinal tract is shunted to the liver where it is deactivated before

30

it can exert its therapeutic effect at the target organ which may be far removed from the liver.

5 The limitations of steroid absorption by both the oral and parenteral routes have lead  
researchers to invent and develop new and more powerful versions of the naturally  
occurring steroids. The resulting compounds exert their desired therapeutic effects at low  
concentration because they are more powerful than the naturally occurring steroids produced  
by the body. However, coupled with these powerful therapeutic effects are frequently  
10 equally powerful side effects. For example the anti-inflammatory corticosteroid prednisone  
is effective in reducing the symptoms associated with many auto-immune diseases such as  
ulcerative colitis, Chron's disease and rheumatoid arthritis, but this powerful steroid also  
causes fluid retention and dramatic loss of bone or osteoporosis.

15 Another result of the limitations of steroid administration is that naturally occurring steroids  
are not utilized effectively as therapeutics. The use of naturally occurring steroids instead of  
the more powerful synthetic compounds is highly desirable because the naturally occurring  
steroids exert their beneficial effects with few if any side effects. Unfortunately, such high  
levels of naturally occurring steroid administration are required to obtain therapeutic benefit  
that the cost of such naturally occurring steroid products becomes prohibitive. Thus, the  
20 possibility of using naturally occurring steroids administered parenterally as therapeutic  
agents has been limited heretofore.

Formulations have been proposed in the past to improve the availability of water-insoluble  
compounds in dosage forms that may be administered parenterally, either intravenously, intra  
25 peritoneally, intrathecally or intramuscularly. Of these routes, the first three are the most  
desirable since the highest concentration of the compound is achieved in the shortest time.

Liposomes, small envelopes of fat-like compounds containing an aqueous chamber or  
chambers within, have been proposed for the parenteral administration of a number of water-  
30 insoluble compounds. The water insoluble compound is dissolved in the fat-like compound  
comprising the envelope of the liposome. Liposomes, however, have the disadvantage of  
being preferentially removed from the circulation and retained in the liver and spleen.

Therefore, unless it is proposed to target the water-insoluble therapeutic compound to these organs, liposomes are not desirable. Liposomes also have a number of other disadvantages, including instability when stored for long and short periods of time. Change in the size of the liposome on storage is also a major problem.

5

Another approach as has been to form a reversible complex between the insoluble drug, such as a steroid, and a carrier molecule. The characteristics of the carrier molecule are such that the carrier molecule and the reversible complex are soluble in water. Among these known carrier molecules are cyclodextrin compounds.

10

A variety of improvements in the characteristics of pharmaceutical complexes including various cyclodextrins and cyclodextrin derivatives are disclosed in the following United States patents:

Noda et al., 4,024,223 methyl salicylate; Szejtli et al 4,228,160 indomethacin; Hyashi et al.,  
15 4,232,009  $\omega$ -halo-PGI<sub>2</sub> analogs; Matsumoto et al., 4,351,846 3-hydroxy and 3-oxo  
prostaglandin analogs; Yamahira et al., 4,353,793, bencyclane fumarate; Lipari, 4,383,992  
steroids-corticosteroids, androgens, anabolic steroids, estrogens, progestagens complexed  
with  $\beta$  cyclodextrin, but not substituted amorphous  $\beta$  cyclodextrins; Nicolau, 4,407,795 P-  
hexadecylaminobenzoic acid sodium salt; Tuttle, 4,424,209 3,4-diisobutyryloxy-N-{3-(4-  
20 isobutyryloxyphenyl)-1-methyl-n-propyl}- $\beta$ -phenethylamine; Tuttle, 4,425,336, 3,4-  
dihydroxy-N-[3-(4-hydroxyphenyl)-1-methyl-n-propyl]- $\beta$ -phenethylamine; Wagu et al.,  
4,438,106 fatty acids EPA and DHA; Masuda et al., 4,474,881 2-(2-fluoro-4-  
biphenyl)propionic acid or salt; Shinoda et al., 4,478,995 acid addition salt of (2'-  
benzyloxycarbonyl)phenyl trans-4-guanidinomehtylcyclo-hexanecarboxylate; Hyashi et al.,  
25 4,479,944 Prostaglandin I<sub>2</sub> analog; Hayashi et al., 4,479,966, 6,9-methano-prostaglandin  
I<sub>2</sub> analogs; Harada et al., 4,497,803 lankacidin-group antibiotic; Masuda 4,499,085  
prostoglandin analog; Szejtli et al., 4,524,068 piperonyl butoxide; Jones, 4,555,504 cardiac  
glycoside; Uekama et al., 4,565,807 pirofen; Ueda et al., 4,575,548 2-nitroxymethyl-6-  
chloropyridine; Ohwaki et al., 4,598,070 triamide anti-hypertensive; Chiesi et al.,  
30 4,603,123 piroxicam (feldene); Hasegawa et al., 4,608,366 monobenzoxamine; Hiari et al.,  
4,659,696 polypeptide; Szejtli et al., 4,623,641 Prostaglandin I<sub>2</sub> methyl ester; Ninger et al.,

4,663,316. unsaturated phosphorous containing antibiotics including phosphotrienin; Fukazawa et al., 4,675,395 hinokitol; Shimizu et al., 4,728,509 3-amino-7-isopropyl-5-oxo--5H-[1]-benzopyrano[2,3-b]pyridine-3-carboxylic acid; Shibani et al., 4,728,510 milk component; and Karl et al., 4,751,095 aspartame.

5

Among the above-mentioned patents, several indicate that complexes of cyclodextrin with drug substances improve side effects of the drug substance. Szejtli et al., 4,228,160 disclosed that the frequency and severity of gastric and duodenal erosion and ulceration in rats caused by indomethacin is improved in an oral formulation of a complex of  $\beta$ -cyclodextrin: indomethacin in a 2:1 ratio, but is not improved and in fact worsens in the same oral formulation of a complex of  $\beta$ -cyclodextrin: indomethacin in a 1:1 ratio.

10

Shimazu et al., 4,352,793 discloses that a formulation wherein bencyclane fumarate an anti-convulsive compound and  $\beta$ -cyclodextrin or  $\gamma$ -cyclodextrin yields a complex in which the bencyclane fumarate is an inclusion compound. These complexes, when formulated as a liquid suitable for oral administration were claimed to be less irritating in an isotonic buffered pH 7 solution when administered as drops to the eyes of rabbits, as compared to bencyclane fumarate drops at the same drug concentration. Shimazu et al., also discloses that similar complexes dissolved in rabbit blood in vitro yielded reduced hemolysis as compared to equal concentrations of bencyclane fumarate alone mixed with rabbit blood.

15

20

Masuda et al., U.S. Patent No. 4,478,811 disclose ophthalmic formulations of  $\beta$ - or  $\gamma$ -cyclodextrin complexes of the nonsteroidal anti-inflammatory compound fluoro-bi-phenylacetic acid which are less irritating and painful than the same formulations of fluoro-bi-phenyl acetic acid alone.

25

Uekama et al., 4,565,807 discloses complexes of  $\alpha$ -,  $\beta$ - and  $\gamma$ - cyclodextrin, pirofen and a pharmaceutically acceptable base. Pirofen is an analgesic and anti-inflammatory compound which is bitter and can cause irritation to the gastrointestinal tract. The complexes disclosed in the patent have improved less bitter taste and are less gastrointestinal irritating than the un-complexed compound pirofen. No preparations suitable for

30

intravenous injection were disclosed.

Lipari, 4,383,992 discloses topical and ophthalmic solutions comprising a number of different steroid-related compounds including corticosteroids, androgens, anabolic steroids, estrogens, and progestagens complexed with  $\beta$  cyclodextrin. None of the cyclodextrin compounds disclosed by Lipari are substituted or amorphous cyclodextrins. In addition, none of the steroid related compounds disclosed by Lipari are  $5\beta$  steroids.

Pitha *et al.*, U.S. Patent No. 4,596,795 discloses complexes containing amorphous hydroxypropyl- $\beta$ -cyclodextrin and sex hormones, particularly testosterone, progesterone and estradiol as a lyophilized powder in tablet form. These tableted complexes are disclosed as appropriate for administration sublingually or buccally with absorption occurring across the corresponding mucosal membrane. None is administered in solution parenterally. In addition, none of the steroid related compounds disclosed by Pitha *et al.* are  $5\beta$  steroids.

Pitha *et al.*, U.S. Patent No. 4,727,064 discloses complexes containing water soluble amorphous substituted cyclodextrin mixtures and drugs with substantially low water solubility which may be lyophilized and the lyophilized powder formed into dosage forms suitable for absorption trans-mucosally across the oral, buccal or rectal mucosa. The solutions of amorphous water soluble cyclodextrin alone and not in a complex with a drug substance are disclosed as nonirritating topically, and having low toxicity, both systemic and local, when applied parenterally. These solutions of substituted cyclodextrin alone were tested and shown to be non-lethal when substantial amounts of the cyclodextrin solution were administered intra peritoneally in mice. There was no disclosure indicating whether the solutions cause elevated temperature in test subjects. This phenomenon called the pyrogenic effect is highly objectionable in parenteral dosage forms. A number of categories of drugs are disclosed in Pitha *et al.* '067 for complex with cyclodextrin derivatives and include *inter alia* vitamins, salts of retinoid acid, spironolactone, antiviral agents, diuretics, anticoagulants, anticonvulsant and anti-inflammatory agents. Significantly, Pitha *et al.* '067 while disclosing that aqueous solutions of 50% cyclodextrin may be used for the purpose of determining solubility of drugs in such solutions does not indicate that such solutions are suitable for intravenous administration. No attempt is made to qualify the solution as to its pyrogenicity.

The claimed compositions of matter in the reference contain only cyclodextrin and drug. Liquid or semi-liquid compositions of matter, which are claimed in the reference, appear to be made of cyclodextrins with higher degrees of substitution with hydroxy propyl groups. These cyclodextrins are themselves semi-solid or liquids according to the reference. Thus no  
5 aqueous formulations of water, cyclodextrin and drug are disclosed or claimed as suitable for parenteral administration in the reference.

Bekers, O., et al., "Stabilization of mitomycins on complex with cyclodextrins in aqueous acidic media" International Journal of Pharmaceutics, 53, 239-248 (1989) describes the  
10 investigation of stabilization of mytomycin-C and several related mitomycins by formation of a complex with cyclodextrin. The authors indicate that at the pH ranges studied  $\alpha$ - and  $\beta$ - cyclodextrin as well as heptakis-(2,6-di-(*l*-methyl)- $\beta$ - cyclodextrin and dimethyl- $\beta$ -cyclodextrin, have no influence on stabilization of mitomycin-C pH degradation.  $\gamma$ -cyclodextrin is reported as having measurable stabilizing effect on mitomycin in acidic media  
15 at pH above 1.

Bodor, US patent 5,024,998 and Bodor, US patent 4,983,586 disclose a series of compositions comprising complexes of hydroxypropyl- $\beta$ -cyclodextrin (HPCD) complexed to a difficult to solubilize drug, or HPCD complexed to a drug which has first been  
20 complexed to a specific class of drug carriers characterized as redox drug carriers. The complex of drug and redox carrier is itself difficult to solubilize and is highly lipophilic due to the presence of pyridine derivatives as part of the redox carrier complex. Bodor '998 and '586 further claim that a solution of 20 to 50% hydroxypropyl- $\beta$ -cyclodextrin and lipophilic drug-redox carrier complex, or 20 to 50% hydroxypropyl- $\beta$ -cyclodextrin and lipophilic  
25 and/or water labile drug is useful in a method of "decreasing the incidence of precipitation of a lipophilic and/or water labile drug occurring at or near the injection site and/or in the lungs or other organs following parenteral administration."

Significantly the Bodor references attribute the precipitation and organ deposition problems  
30 associated with parenteral administration of lipophilic drugs to the effects of organic solvents used to solubilized the drug in the parenteral vehicle. The Bodor references additionally



state that drugs which are particularly useful in the parenteral composition and methods disclosed therein are those which are relatively insoluble in water but whose water solubility can be substantially improved by formulation with 20 to 50% of the selected cyclodextrin, e.g., HPCD, in water. Significantly no part of Bodor addresses the pyrogenic load on the  
5 cyclodextrin or the issue of the pyrogenic effect of the composition when injected parenterally. Thus it is quite clear that the Bodor references are directed to prevention of the phenomenon of precipitation of insoluble drugs and insoluble drug-carrier complexes.

There are a number of disclosures relating to the 5  $\beta$  steroids used in the formulations disclosed herein below according to the invention. Yen *et al.*, *Lipids*, 12, 409 (1977)  
10 disclosed that dehydroepiandrosterone DHEA ( a 5  $\alpha$  steroid) administered by a variety of routes decreased the rate of weight gain in a strain of genetically obese mice. DHEA treatment markedly reduced the development of diabetes in both genetically obese and diabetic mice and maximal benefit was observed when DHEA was ingested according to  
15 Coleman *et al.*, *Diabetes*, 31:80 (1982). Coleman *et al.*, *Endocrinology*, 115, 239 (1984) showed that  $\alpha$  ET and  $\beta$  ET reduce blood sugar, increased plasma insulin concentrations and provided a protective effect on the pancreas as demonstrated by increased granulation of islet  $\beta$  cells. Moreover,  $\alpha$  ET and  $\beta$  ET but not androsterone or epiandrosterone, were four times more effective than DHEA in preventing development of diabetes in C57BL/KsJ-db/db  
20 diabetic mice. Coleman *et al.*, U.S. Patent No. 4,518,595 showed that oral administration of DHEA restored hyperglycemia to normal levels and improved glucose tolerance even in severely diabetic mammals. In U.S. Patent No. 4, 507,289 Coleman taught the use of  $\alpha$  ET and  $\beta$  ET and an estrogen for the treatment of diabetes, obesity syndromes and associated hypercorticism.

25 Coleman, *Endocrinology*, 117, 2279 (1985) disclosed that  $\alpha$  ET and  $\beta$  ET when supplied in the diet have anti-obesity properties, and can prevent and arrest the development of obesity, and facilitate weight reduction after obesity in diabetic genetically obese mice. U.S. Patent No. 4,666,898 to Coleman and Applezweig disclosed the use of Etiocholanolones in  
30 the treatment of obesity, diabetes and other symptoms of hypercorticism. B. Zumoff *et al.*, *Obesity Research*, 2, 13 (1994) disclosed that ED orally administered at a dose of four

grams per day yielded significant fat loss in human obese subjects. In a 20 week randomized double-blind cross over study, 14 subjects lost significantly more weight and body fat during treatment with oral ED than during placebo administration. Mean weight loss during ED administration was  $2.8 \pm 5.5$  kilograms which was equivalent to  $0.5 \pm 0.91$  kilograms per week per 100 kilograms of body fat. Densitometric measurement of body fat content showed that the mean weight loss coincided almost exactly with the mean decrease in body fat content. Over the 10-week period of ED administration, the mean fat loss was about 5% of the initial body fat content. There were no significant subjective or objective side effects of ED administration. In all of the foregoing references, the  $5\beta$  steroids or DHEA is administered orally with or without food.

U.S. Patent 5,006,517 to Bradlow, *et al.*, discloses that Prader-Willi Syndrome a congenital disease caused by a chromosomal defect may be treated by administering etiocholanolone or etiocholanolondione to individuals suffering from the syndrome resulting in either weight loss or a decrease in the rate of weight gain in the treated individuals.

Gardner and Juneja, *Brit. J. Hemat.*, 65,295-300 (1987) treated aplastic anemia patients with either  $\alpha$ - and/or  $\beta$  ET in an uncontrolled pilot study. Of 43 treated patients, 17 had hematologic responses lasting longer than six months including normalization of hemoglobin level for long periods of time which persisted after  $\alpha$ -ET was discontinued. Kappas *et al.*, *J. Clin Endocr.*, 16, 284 (1956); Kappas *et al.*, *Trans. Assn. Am. Phys.*, 72,54 (1959) showed that  $\alpha$ -ET engendered acute pyrogenic reactions in man when administered by intramuscular injection. This observation lead Gardner and Juneja to concomitantly administer prednisolone along with the intramuscular therapeutic injections of  $\alpha$ -ET to prevent development of fever and to alleviate local irritation at the injection sites. The  $\alpha$ -ET and  $\beta$ -ET as described below, were administered in a sterile polypropylene vehicle by im route because the  $5\beta$  steroids are notoriously insoluble in aqueous solutions and therefore cannot be administered by intravenous route.

Gardner and Juneja also observed that three patients who did not respond to  $\alpha$  ET had a

hematologic recovery when treated with  $\beta$ -ET. Significantly,  $\beta$ -ET evidenced little pyrogenic effect and minimal prednisolone doses of 10 mg were sufficient to suppress any local irritation.

5 The metabolism and elimination routes of the 5  $\beta$  steroids have been investigated by several groups. Administration of  $\alpha$  ET and  $\beta$  ET by parenteral routes resulted in the recovery of almost exclusively  $\alpha$  ET conjugates in the urine, Kappas *et al.*, J. Clin Endocr., 16, 284 (1956). In addition Zumoff *et al.*, have shown that  $\beta$  ET or ED when administered to a patient is quickly converted almost exclusively to  $\alpha$  ET in the serum of normal and obese  
10 subjects (unpublished data). Bradlow *et al.*, J. Clin Endocr. 27,1203-1207 (1967) found that based on isotopic ratio studies, the C3 position of a portion of  $\alpha$  ET administered IV to human subjects was rapidly oxidized and re-reduced.

Appelzweig *et al.*, U.S. Patent No. 4,871,726 disclosed methods and compositions for  
15 inducing increased blood levels of  $\alpha$ -ET in the body of a mammal by administration of ED, showing that both  $\alpha$  ET and  $\beta$  ET administered either orally or parenterally are rapidly oxidized at the C-3 position to form etiocholandione (ED). Furthermore they disclose that ED can serve as a source of circulating blood levels of free  $\alpha$  ET, By administering ED, high levels of  $\alpha$  ET can be achieved in the serum. In effect ED can serve as a prodrug for  $\alpha$  ET  
20 according to Appelzweig *et al.*

While several of the 5  $\beta$  steroids are known to be active in control of obesity and associated diabetic and/or hypercortical syndrome, the compounds are effective for treatment of these conditions when they are administered orally. By the oral route, the absorption of the 5 $\beta$   
25 steroids is only 5 to 15% as measured by blood levels using various assays. Thus a large portion of the administered drug is never absorbed into the blood stream and the greatest part of the drug that is administered is eliminated in the feces.

To achieve high serum concentrations of a drug, it is more efficient to administer drugs  
30 parenterally, preferably by the intravenous (iv) route. However, it has been observed that the 5 $\beta$  steroids do not appear to exert any of their anti-obesity, anti-diabetic or anti-hypercortical

activities when administered intramuscularly (im). Furthermore the im route of administration engenders pyrogenic reactions with some of the  $5\beta$  steroids as described above in Kappas *et al.* Thus, it would be desirable to have a formulation that would allow parenteral administration of the  $5\beta$  steroids, preferably by the intravenous (iv) route and  
5 which would preserve the anti-obesity, anti-diabetic and anti-hypercortical activities of these compounds.

A further long-felt need in the administration of these  $5\beta$  steroid drugs is to obtain a high serum concentration of the drug without resorting to the intramuscular (im) route of  
10 administration which causes local irritation and, at least in the case of  $\alpha$ -ET pyrogenic reaction and notable patient discomfort. Furthermore, in that the evidence of Gardner and Juneja implies that pyrogenic reaction associated with im administration is not required to obtain the hematologic activity of the  $5\beta$  steroids, especially  $\beta$ -ET, it would be desirable to obtain a formulation of the  $5\beta$  steroids compatible with aqueous iv solutions to maximize the  
15 serum concentration of the  $5\beta$  steroids. Considerable saving in the cost of active drug substance for a given response could be achieved thereby.

In addition to the advantages of higher serum concentration of drug and drug precursors afforded by a practical aqueous-compatible formulation of  $\alpha$  ET and ED respectively, it  
20 would be of considerable advantage to provide an aqueous compatible formulation of the drug and prodrug which further includes an amount of  $\beta$  ET which would be clinically effective especially in anemia patients who respond hematologically to  $\beta$  ET but who do not respond to  $\alpha$  ET.

25 The steroid dehydroepiandrosterone (3- $\beta$ -hydroxy-androst-5-en-17-one, DHEA) and its sulfate derivatives are major steroid adrenal secretory products in humans. DHEA is metabolized to testosterone (17- $\beta$ -hydroxy-androst-4-en-3-one) and estradiol (estra-1, 3,5 (10)-triene-3, 17-diol), two major sex hormones in humans. Other metabolites of DHEA include  $\alpha$  ET and  $\beta$ -ET. They were considered to be inert metabolic end products which  
30 were merely conjugated as glucuronides or sulfates and excreted into the urine.  $\alpha$  ET is a major metabolite of DHEA, and in normal individuals, is excreted in the urine in amounts of

about 3-5 mg per day, whereas  $\beta$  ET is a minor metabolite in man.

DHEA has been shown to be effective in controlling diabetes and obesity in rats and mice by Coleman, however in human usage for a number of different inflammatory conditions including the treatment of Lupus Erythematosus, DHEA has demonstrated a number of undesirable side effects. In particular, in female patients, who are the most frequent sufferers of lupus erythematosus, DHEA causes severe acne and may also cause masculinizing effects, particularly hirsutism, the production of facial and body hair.

One of the advantages that may be obtained through the present invention is to reduce the amount of DHEA administered to a patient when administered in an iv formulation. The administration of DHEA by the iv route is advantageous in itself in that the circulating concentrations of the drug can be precisely monitored. If orally administered, the amount of DHEA absorbed through the alimentary tract and reaching the blood stream may vary substantially depending upon the amount of food and liquid the subject has consumed. Furthermore since many inflammatory processes including Chron's disease and ulcerative colitis affect the intestines and bowel, the absorption of a drug from this area of the body may vary considerably if orally administered. Thus, iv administration may be particularly important for DHEA.

In particular it is believed that some of the salutary effects of DHEA, particularly those relating to inflammatory and autoimmune processes may be preserved while the side effects are eliminated if reduced amounts of DHEA are administered by the parenteral route, especially the iv route. In addition, it is believed that the salutary effects of DHEA on inflammatory and autoimmune disease, diabetes and obesity may be preserved even at low concentrations if DHEA is co-administered with one or more of the  $5\beta$  steroids parenterally. The parenteral administration of DHEA with one or more of the  $5\beta$  steroids is facilitated in the formulation according to the invention.

Summary and Objects of the Invention

It is an object of the invention to provide improved formulations of the 5  $\beta$  steroids suitable for parenteral administration.

5 It is a further object of the invention to provide improved formulations of the 5  $\beta$  steroids wherein the formulation is a sterile aqueous solution suitable for parenteral administration.

It is yet a further object of the invention to provide improved formulations of the 5  $\beta$  steroids wherein the formulation is a sterile aqueous solution suitable for parenteral administration and is not pyrogenic.

10

It is a still further object of the invention to provide improved formulations of the 5  $\beta$  steroids wherein the formulation is non-pyrogenic and contains one or more of the 5  $\beta$  steroids.

15 It is yet still a further object of the invention to provide improved formulations of the 5  $\beta$  steroids wherein the formulation is non-pyrogenic and contains one or more of the 5  $\beta$  steroids and DHEA.

Another object of the invention is to provide methods for treating a subject for a condition which responds to treatment with the 5  $\beta$  steroids by administering to the subject an improved formulation of the 5  $\beta$  steroids suitable for parenteral administration. Among these conditions may be obesity, diabetes syndrome, diabetes-associated hypercorticism or combinations thereof, anemic disorders including but not limited to aplastic anemia and anemia associated with renal failure or chemotherapy-induced or radiation-induced anemia or neutropenia, autoimmune or inflammatory disorders such as lupus erythematosus.

25

It is a further object of the invention to provide a method for treating a subject for a condition which responds to treatment with DHEA using a reduced amount of DHEA by administering to the subject at least one 5  $\beta$  steroid and a reduced amount of DHEA. Among these conditions are those listed above.

30

Yet a still further object of the invention is to provide a method for treating a subject for a

condition which responds to treatment with DHEA using a reduced amount of DHEA by administering to the subject at least one 5  $\beta$  steroid and a reduced amount of DHEA in a formulation including an amorphous cyclodextrin wherein the formulation is suitable for parenteral administration. Among these conditions are those listed above.

5

Another object of the invention is to provide a method of treating a subject for a condition which responds to treatment with the 5  $\beta$  steroids by administering to the subject an improved formulation of the 5  $\beta$  steroids suitable for parenteral administration which is non-pyrogenic. Among these conditions are those listed above.

10

#### Detailed Description of the Invention

15

By 5  $\beta$  steroid is meant  $\alpha$  ET,  $\beta$  ET, and ED. In addition certain alkylated derivatives of these 5  $\beta$  steroids are also included in this definition. For example 16-alkylated 5 $\beta$  androstan-3-ol-17-one and 16-alkylated 5  $\beta$  androstan-3, 7-diol-17-one which are disclosed in U.S. Patent 4,602,008, herein incorporated by reference, have been shown to be biologically effective as anti-diabetic, anti-obesity and erythropoietic agents in mammals. These compounds may be esterified at the 3 or 7 positions of the steroid ring structure by conventional means. Also included in the definition is 5- $\beta$ -androstan-3,17-diol or etiocholandiols (herein after Ediol.)  $\alpha$  ET,  $\beta$  ET, Ediol and ED are all commercially available compounds (Research Plus, Inc, POB 324, Bayonne, New Jersey 07002 USA).

25

In addition other diols of the 17 keto 5- $\beta$  steroids are commercially available and may be used in the invention and may be esterified at the carbon at which the -OH moiety is attached. Thus, for example 5 $\beta$ -androstan-3 $\alpha$ , 11 $\alpha$ -diol-17-one, 5 $\beta$ -androstan-3 $\beta$ , 11 $\beta$ -diol-17-one, and 5 $\beta$ -androstan-3 $\alpha$ , 11 $\beta$ -diol-17-one, 5 $\beta$ -androstan-11 $\alpha$ -ol-3, 17-dione, 5 $\beta$ -androstan-11 $\beta$ -ol-3, 17-dione, 5 $\beta$ -androstan-3 $\alpha$ , 16 $\alpha$ -diol-17-one are commercially available

30

(Research Plus, Inc, POB 324, Bayonne, New Jersey 07002 USA) and may be esterified at the 11 or 16 positions, as the case may be, by conventional means to produce the organic acid derivative of the 5 $\beta$  steroids.

5 By cyclodextrin is meant  $\alpha$ -,  $\beta$ -, or  $\gamma$ - cyclodextrin. Cyclodextrins are described in detail in Pitha et al., U. S. Patent 4,727,064 which is incorporated herein by reference. Cyclodextrins are cyclic oligomers of glucose; these compounds form inclusion complexes with any drug whose molecule can fit into the lipophile-seeking cavities of the cyclodextrin molecule.

10

By amorphous cyclodextrin is meant non-crystalline mixtures of cyclodextrins wherein the mixture is prepared from  $\alpha$ -,  $\beta$ -, or  $\gamma$ - cyclodextrin. In general the amorphous cyclodextrin is prepared by non-selective additions, especially alkylation of the desired cyclodextrin species. Reactions are carried out to yield mixtures containing a plurality of  
15 components thereby preventing crystallization of the cyclodextrin. various alkylated and hydroxyalkyl-cyclodextrins can be made and of course will vary, depending upon the starting species of cyclodextrin and the addition agent used. Among the amorphous cyclodextrins suitable for compositions according to the invention are hydroxypropyl, hydroxyethyl, glucosyl, maltosyl and maltotriosyl derivatives of  $\beta$ -cyclodextrin, carboxyamidomethyl- $\beta$ -  
20 cyclodextrin, carboxymethyl- $\beta$ -cyclodextrin, hydroxypropyl- $\beta$ -cyclodextrin and diethylamino- $\beta$ -cyclodextrin. In the compositions according to the invention hydroxy- $\beta$ -cyclodextrin is preferred. The substituted  $\gamma$ - cyclodextrins may also be suitable, including hydroxypropyl, hydroxyethyl, glucosyl, maltosyl and maltotriosyl derivatives of  $\gamma$ -cyclodextrin.

25

The cyclodextrin of the compositions according to the invention may be  $\alpha$ -,  $\beta$ -, or  $\gamma$ -cyclodextrin.  $\alpha$ -cyclodextrin contains six glucopyranose units;  $\beta$ -cyclodextrin contains seven glucopyranose units; and  $\gamma$ -cyclodextrin contains eight glucopyranose units. The  
30 molecule is believed to form a truncated cone having a core opening of 4.7-5.3 Å, 6.0-6.5 Å and 7.5-8.3 Å in  $\alpha$ -,  $\beta$ -, or  $\gamma$ - cyclodextrin respectively. The composition according to



the invention may comprise a mixture of two or more of the  $\alpha$ -,  $\beta$ -, or  $\gamma$ - cyclodextrins. Usually, however the composition according to the invention will comprise only one of the  $\alpha$ -,  $\beta$ -, or  $\gamma$ - cyclodextrins. The particular  $\alpha$ -,  $\beta$ -, or  $\gamma$ - cyclodextrin to be used with the particular  $5\beta$  steroid compound to form the compositions according to the invention may  
5 be selected based on the known size of the molecule of the  $5\beta$  steroid compound and the relative size of the cavity of the cyclodextrin compound. Generally if the molecule of the  $5\beta$  steroid compound is relatively large, a cyclodextrin having a larger cavity is used to make the composition according to the invention. Furthermore, if the  $5\beta$  steroid compound is administered with an excipient it may be desirable to use a cyclodextrin compound having  
10 a larger cavity in the composition according to the invention.

The unmodified  $\alpha$ -,  $\beta$ -, or  $\gamma$ - cyclodextrins are less preferred in the compositions according to the invention because the unmodified forms tend to crystallize and are relatively less soluble in aqueous solutions. More preferred for the compositions according to the  
15 invention are the  $\alpha$ -,  $\beta$ -, and  $\gamma$ - cyclodextrins that are chemically modified or substituted. Chemical substitution at the 2,3 and 6 hydroxyl groups of the glucopyranose units of the cyclodextrin rings yields increases in solubility of the cyclodextrin compound.

Most preferred cyclodextrins in the compositions according to the invention are amorphous  
20 cyclodextrin compounds. By amorphous cyclodextrin is meant non-crystalline mixtures of cyclodextrins wherein the mixture is prepared from  $\alpha$ -,  $\beta$ -, or  $\gamma$ - cyclodextrin. In general, the amorphous cyclodextrin is prepared by non-selective alkylation of the desired cyclodextrin species. Suitable alkylation agents for this purpose include but are not limited to propylene oxide, glycidol, iodoacetamide, chloroacetate, and 2-  
25 diethylaminoethylchloride. Reactions are carried out to yield mixtures containing a plurality of components thereby preventing crystallization of the cyclodextrin. Various alkylated cyclodextrins can be made and of course will vary, depending upon the starting species of cyclodextrin and the alkylating agent used. Among the amorphous cyclodextrins suitable for compositions according to the invention are hydroxypropyl, hydroxyethyl, glucosyl, maltosyl  
30 and maltotriosyl derivatives of  $\beta$ -cyclodextrin, carboxyamidomethyl- $\beta$ -cyclodextrin, carboxymethyl- $\beta$ -cyclodextrin, hydroxypropyl- $\beta$ -cyclodextrin and diethylamino- $\beta$ -

cyclodextrin. In the compositions according to the invention hydroxypropyl- $\beta$ -cyclodextrin is preferred although the  $\alpha$ - or  $\gamma$ - analogs may also be suitable. The particular alkylated  $\alpha$ -,  $\beta$ -, or  $\gamma$ - cyclodextrin to be used with the particular 5  $\beta$  steroid compound to form the compositions according to the invention will be selected based on the size of the molecule of the 5  $\beta$  steroid compound and the relative size of the cavity of the cyclodextrin compound.

As with the unsubstituted cyclodextrins mentioned above, it may be advantageous to use alkylated cyclodextrin having a larger cavity when the composition according to the invention also includes an excipient. The use of a particular  $\alpha$ -,  $\beta$ -, or  $\gamma$ - cyclodextrin with a particular 5  $\beta$  steroid compound or 5  $\beta$  steroid compound and excipient in the compositions according to the invention may of course be optimized based on the effectiveness in of maintaining the 5  $\beta$  steroid or mixture thereof with or without DHEA in solution.

As mentioned above, the compositions of matter of the invention comprise an aqueous preparation of preferably substituted amorphous cyclodextrin and one or more 5  $\beta$  steroids.

The relative amounts of 5  $\beta$  steroid compound and cyclodextrin will vary depending upon the relative amount of each of the 5  $\beta$  steroid compounds and the effect of the cyclodextrin on the compound. In general, the ratio of the weight of 5  $\beta$  steroid compound to the weight of cyclodextrin compound will be in a range between 1:1 and 1:5000. Within this range, the circulating availability of the 5  $\beta$  steroid will be significantly increased when the ratio of the weight of 5  $\beta$  steroid to the weight of cyclodextrin compound is in a range between the concentration at which the 5  $\beta$  steroid will not go into solution at the particular amorphous cyclodextrin concentration and 1:2000. A weight to weight ratio in a range of 1:5 to 1:200 and more preferably in a range of 1:5 to 1:50 of 5  $\beta$  steroid compound to cyclodextrin are believed to be the most effective for increased circulating availability of the 5  $\beta$  steroid. For example, ED in a ratio of between 1:10 and 1:300 (drug: amorphous cyclodextrin, wt:wt), and a final concentration of the injection solution of 40 mg/ml of ED is expected to significantly increase circulating concentration of ED as compared to intramuscularly injected ED in a polypropylene vehicle.

Thus  $\alpha$  ET,  $\beta$  ET, ED, Ediol or DHEA each individually or in a mixture of two or more may be added to the cyclodextrin solution to form a complex of each compound individually

or in a mixture of two or more of these compounds.

Importantly, if the aqueous solution comprising the DHEA and /or 5  $\beta$  steroid and amorphous cyclodextrin is to be administered parenterally, especially via the iv route, the amorphous cyclodextrin will be substantially free of pyrogenic contaminants. Amorphous hydroxypropyl- $\beta$ -cyclodextrin maybe purchased from a number of vendors including Amaizo, Inc. (Hammond Indiana, USA) under the tradename Encapsin. In addition, other forms of amorphous cyclodextrin having different degrees of substitution or glucose residue number are available commercially. A method for the production of hydroxypropyl- $\beta$ -cyclodextrin is disclosed in Pitha *et al.*, U.S. Patent No.4, 727,064 which is incorporated herein by reference.

To produce the formulations according to the invention, a pre-weighed amount of hydroxypropyl- $\beta$ -cyclodextrin compound, which is substantially pyrogen free is placed in a suitable depyrogenated sterile container. Methods for depyrogenation of containers and closure components are well known to those skilled in the art and are fully described in the United States Pharmacopeia 23 (United States Pharmacopeial Convention, Rockville, Maryland USA). Generally, depyrogenation is accomplished by exposing the objects to be depyrogenated to temperatures above 400°C for a period of time sufficient to fully incinerate any organic matter. As measured in U.S.P. Bacterial Endotoxin Units, the formulation will contain no more than 10 Bacterial Endotoxin Units per gram of amorphous cyclodextrin. By substantially pyrogen free is meant that the hydroxypropyl- $\beta$ -cyclodextrin contains less than 10 U.S.P. bacterial endotoxin units per gram using the U.S.P. method. Preferably, the hydroxypropyl- $\beta$ -cyclodextrin will contain between 0.1 and 5 U.S.P. bacterial endotoxin units per mg, under conditions specified in the United States Pharmacopeia 23.

Sufficient sterile water for injection is added to the substantially pyrogen free amorphous cyclodextrin until the desired concentration of hydroxypropyl- $\beta$ -cyclodextrin is in solution. To this solution a pre-weighed amount of the 5  $\beta$  steroid is added with agitation and with additional standing if necessary until it dissolves.

The solution is then filtered through a sterile 0.2 micron filter into a sterile holding vessel and is subsequently filled in sterile depyrogenated vials and is capped. For products that will be stored for long periods of time, a pharmaceutically acceptable preservative may be added to the solution of  $5\beta$  steroid and hydroxypropyl- $\beta$ -cyclodextrin prior to filtration, filling and capping or alternatively, may be added sterilely after filtration.

In the formulations according to the invention, DHEA alone or together with one or more of the  $5\beta$  steroids may be used. In particular the formulation including DHEA and ED or DHEA and  $\alpha$ ET and  $\beta$ ET or DHEA, ED,  $\alpha$ ET and  $\beta$ ET are desirable. This is particularly desirable in relation to obesity control and inflammatory disease control. It is clear that DHEA is metabolically transformed into the primary male and female sex steroid hormones, testosterone and estradiol respectively, and is ultimately metabolized to  $\alpha$  ET and  $\beta$  ET. By reducing the amount of DHEA to an amount which contributes in only a small way to the production of the sex steroids testosterone and estradiol, while at the same time maximizing the amount of  $\alpha$  ET and  $\beta$ ET circulating in the blood from the initial dose administered to a subject, it is believed that enhancement of the desirable obesity control, hematopoiesis, and anti-inflammatory effects may be achieved while, at the same time, the dermatologic and virilizing side effects of DHEA are minimized to an acceptable degree, if not eliminated.

20

The invention will be better understood from the following examples which are intended to be merely illustrative of the invention and are not intended to be limiting.

#### Examples

##### Example I

12 milligrams (mg) of Etiocholandione (Steraloids, Inc. New Hampshire) was stirred and shaken with 0.5 ml of water in a test tube. Appreciable quantities of compound remained out of solution after 5 minutes accumulating as white crystals at the bottom of the test tube.

A 50% solution of hydroxypropyl- $\beta$ -cyclodextrin was prepared as follows. 5 grams of pyrogen free hydroxypropyl- $\beta$ -cyclodextrin (sold under the trade name Encapsin, Amaizo, Inc., Hammond, Indiana, USA) was weighed on an analytical scale and placed in a graduated

cylinder. Water was added with shaking until the volume reached 10 ml.

0.5 ml of the 50% solution of hydroxypropyl- $\beta$ -cyclodextrin was added to the 0.5 ml of water containing 12 mg of Etiocholandione with additional shaking. A clear solution was obtained.

5 Thus, 12 mg ED was effectively solubilized in 1 ml of 25% solution of hydroxypropyl- $\beta$ -cyclodextrin.

#### Example II

10 The previous experiment was repeated using a 50% solution of hydroxypropyl- $\beta$ -cyclodextrin prepared as in Example I. 50 mg of Etiocholandione was weighed and placed in a test tube. 2 ml of the 50% solution of hydroxypropyl- $\beta$ -cyclodextrin was added to the test tube with shaking. Within one minute, the Etiocholandione dissolved, leaving a clear solution with no accumulation of crystals. Thus, 25 mg was effectively solubilized in 1 ml of 50% solution  
15 of hydroxypropyl- $\beta$ -cyclodextrin.

#### Example III

4 grams of hydroxypropyl- $\beta$ -cyclodextrin was dissolved in 7 ml of distilled water to a total  
20 volume of 10 ml. Each ml of stock solution contained 0.4 g of solution hydroxypropyl- $\beta$ -cyclodextrin. 20 mg of 3- $\beta$ -hydroxy androstan-17-one ( $\beta$ -etiocholanolone) was shaken with 1 ml of the freshly prepared stock solution. Partial dissolution took place immediately and after sitting at room temperature with occasional shaking for 6 hours complete dissolution took place.

25

#### Example IV

The same experiment as in example III was performed except that 40 mg of  $\beta$ -etiocholanolone was added to the stock solution of hydroxypropyl- $\beta$ -cyclodextrin. After  
30 standing for 24 hr, a complete dissolution had occurred.

#### Example V

The same experiment as in example III was performed except that 20 mg of Etiocholandione was added to the stock solution of hydroxypropyl- $\beta$ -cyclodextrin. Complete dissolution  
5 occurred within 6 hours.

#### Example VI

The same experiment as in example V was performed except that 40 mg of Etiocholandione  
10 was added to the stock solution of hydroxypropyl- $\beta$ -cyclodextrin. Complete dissolution occurred within 24 hours.

#### Example VII

15 The same experiment as in Example V was performed except 80 mg or 250mg ED was added to the stock solution. In neither case did the ED completely dissolve.

#### Example VII

20

The optical density (OD) of the ET and ED dissolved in hydroxypropyl- $\beta$ -cyclodextrin solutions was determined at 400 NM against a blank control containing water and set at an OD of 0.0. The dissolved solutions of ET and ED had OD readings of 0.12 whereas the hydroxypropyl- $\beta$ -cyclodextrin solution containing 80mg of ED gave an OD of 1.2 at 499  
25 NM.

#### Example VIII

30 To make a formulation containing  $\alpha$  ET and ED, the stock solution of hydroxypropyl- $\beta$ -

cyclodextrin in prepared as in Example III. 20 mg of  $\alpha$  ET and 20 mg of ED will be added to 1 ml of the solution with agitation. An aqueous solution containing 40 mg of the two 5  $\beta$  steriods and no precipitate is present after standing for 24 hours.

5 Example IX

To make a formulation containing  $\beta$  ET and ED, the stock solution of hydroxypropyl- $\beta$ -cyclodextrin in prepared as in Example III. 20 mg of  $\beta$  ET and 20 mg of ED is added to 1 ml of the solution with agitation. An aqueous solution containing 40 mg of the two 5  $\beta$  steriods and no precipitate is present after standing for 24 hours.

Example X

To make a formulation containing  $\alpha$  ET,  $\beta$  ET and ED, the stock solution of hydroxypropyl- $\beta$ -cyclodextrin is prepared as in Example III. 15 mg of  $\alpha$  ET, 15 mg  $\beta$  ET and 15 mg of ED will be added to 1 ml of the solution with agitation. A clear aqueous solution containing 45 mg of the three 5  $\beta$  steriods and no precipitate will be present after standing for 24 hours.

Example XI

20

To make a formulation containing  $\alpha$  ET,  $\beta$  ET and ED, the stock solution of hydroxypropyl- $\beta$ -cyclodextrin is prepared as in Example III. 10 mg of  $\alpha$  ET, 10 mg  $\beta$  ET, 10 mg of ED and 15 mg of DHEA will be added to 1 ml of the solution with agitation. A clear aqueous solution containing 45 mg of the DHEA and the three 5  $\beta$  steriods and no precipitate is present after standing for 24 hours.

25

The composition of matter according to the invention may be supplied as a dry powder or as a solution. If the composition of matter is to be injected into a subject it will be rendered sterile prior to injection. Accordingly, the composition of matter according to the invention may be supplied as a sterile cake, plug or powder or as a sterile lyophilized preparation in a

30

sterile vial suitable for the addition of a sterile diluent, or as a sterile liquid solution in a sterile container.

5 The compositions of matter according to the invention may be supplied as a powder comprising the active pharmaceutical 5  $\beta$  steroid compound or compounds and amorphous cyclodextrin compound. If the composition is to be administered parenterally, for example iv, the composition of matter will be rendered sterile prior to such administration. Any of the several known means for rendering such pharmaceutical preparations sterile may be used so long as the active pharmaceutical compound is not inactivated and the complex with the  
10 amorphous cyclodextrin is not degraded. If the active pharmaceutical compound is heat stable, the composition of matter according to the invention may be heat sterilized. If the cytotoxic compound is not heat-stable but is not photo degraded the composition may be sterilized by exposure to ultraviolet light or by ionizing radiation. Alternatively, the composition of matter if in a powder form may be gas sterilized using for example ethylene  
15 oxide gas. In another alternative, the composition of matter according to the invention may be filter-sterilized using a 0.2 micron filter. If the composition of matter is an aqueous liquid, it may be filled in a sterile container and supplied as a sterile liquid ready for further dilution or injection neat. Alternatively such sterile liquids may be freeze-dried or lyophilized in a sterile container and capped.

20

In general the compositions of matter according to the invention will be made by dissolving the cyclodextrin in water and adding the 5 $\beta$  steroid compound to the aqueous cyclodextrin solution. Excipients, if any are desired, may be added with or subsequent to adding the 5 $\beta$  steroid compound. The resulting solution may be sterilized using any of the known methods  
25 appropriate to preserving the compound without significant degradation.

Preferably the solution will be sterile filtered, although other means such as terminal heat sterilization or irradiation may be employed as is known in the art, provided that the cyclodextrin compound is not significantly degraded. Alternatively, the components may be  
30 sterilized by any of the known methods appropriate to preserving the compound prior to



mixing in water and may be mixed using sterile equipment and technique. The solution may be lyophilized in sterile containers and capped. Prior to use the lyophilized composition of matter may be reconstituted using sterile water for injection.

- 5 The container closure system used for containing the formulation according to the invention will also be treated to remove or destroy pyrogenic substances by means known in the art prior to filling and further processing. Thus the preferred compositions of matter according to the invention for parenteral administration, especially by the intravenous route will be non-pyrogenic. Nonpyrogenic preparations according to the invention, when administered to a  
10 subject, does not cause a febrile (basal body temperature raising) reaction. Although some bacterial endotoxin may be present, the amount is insufficient to elicit a febrile reaction. In general, such non-pyrogenic compositions will contain less than 10 U.S.P. bacterial endotoxin units per gram of product.
- 15 The formulation according to the invention may be supplied as a dry lyophilized powder as mentioned above or as a sterile non pyrogenic aqueous solution in a sterile container closure system such as a stoppered vial suitable for puncturing with a sterile syringe and needle.

Alternatively the formulation according to the invention may be supplied as a sterile non  
20 pyrogenic aqueous solution in a sterile syringe or syringe and needle. As a sterile solution or powder it may also include a pharmaceutically acceptable preservative. The formulation according to the invention may also be included in other dosage forms in addition to those appropriate for parenteral administration. Preferably, such other dosage forms will include one or more of the 5  $\beta$  steroids with or without DHEA. Such dosage forms may be in the  
25 form of aqueous suspensions, elixirs, or syrups suitable for oral administration, or compounded as a cream or ointment in a pharmaceutically acceptable topical base allowing the 5  $\beta$  steroid compounds with or without DHEA to be absorbed across the skin. In addition the formulation according to the invention may be compounded in a lozenge or suppository suitable for trans mucosal absorption.

30

The formulations according to the invention having been described herein may influence the ordinarily skilled artisan to make similar formulations using components that will be known in the art, without departing from the invention which is claimed herein.

We Claim:

- 5 1. A composition of matter comprising at least one  $\Delta^5$  steroid and an amorphous cyclodextrin.
2. The composition of claim 1 wherein said  $\Delta^5$  steroid is selected from the group consisting  
10 of  $\alpha$  etiocholanolone,  $\beta$  etiocholanolone and etiocholandione.
3. The composition of claim 1 comprising etiocholandione and  $\beta$  etiocholanolone.
4. The composition of claim 1 comprising etiocholandione and  $\alpha$  etiocholanolone.  
15 5. The composition of claim 1 comprising etiocholandione,  $\alpha$  etiocholanolone and  $\beta$  etiocholanolone.
6. A composition wherein the any of the forgoing compositions is a sterile aqueous solution  
20 suitable for parenteral administration.
7. The composition of claim 6 wherein the composition is sterile and is contained within a sterile container.
8. The composition of claim 7 wherein the composition is non-pyrogenic when administered  
25 intravenously.
9. The composition of claim 8 wherein said composition contains no more than 10 endotoxin units per gram.
10. The composition wherein in any of the forgoing compositions said amorphous cyclodextrin has a degree of substitution of 2 to 7.

30

11. The composition wherein any of the forgoing compositions further includes DHEA.
12. A composition of any of the forgoing claims suitable for treating a condition selected from the group consisting of obesity, diabetes syndrome, diabetes-associated  
5 hypercorticism, combinations thereof, and anemic disorders wherein the  $5\beta$  steroid or  $5\beta$  steroids together is an obesity-, diabetes-, hypercorticism- or anemia- antagonistic amount.

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/US 96/18189

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 6 A61K47/40 A61K47/48 A61K31/565

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)  
IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 477 107 A (LAB. BESINS ISCOVESCO SA) 25 March 1992 see page 3, line 4 - line 7 see page 9; table 2 ---	1,2,6-10
X	JOURNAL OF PHARMACEUTICAL SCIENCES, vol. 84, no. 10, 1 October 1995, pages 1154-1159, XP000529726 BREWSTER M E ET AL: "PREPARATION, CHARACTERIZATION, AND ANESTHETIC PROPERTIES OF 2-HYDROXYPROPYL-B-CYCLODEXTRIN COMPLEXES OF PREGNANOLONE AND PREGNENOLONE IN RAT AND MOUSE" see the whole document ---	1,10
A	WO 92 04888 A (US ARMY) 2 April 1992 see page 13; example 1 ---	1
-/--		

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*&\* document member of the same patent family

2

Date of the actual completion of the international search  1 April 1997	Date of mailing of the international search report  16.04.97
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+ 31-70) 340-3016	Authorized officer  Boulois, D

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 96/18189

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0 149 220 A (JACKSON LAB ;PROGENICS INC (US)) 24 July 1985 cited in the application see the whole document ---	12
A	EP 0 261 891 A (PROGENICS INC) 30 March 1988 see the whole document ---	12
X	J. OF LIQUID CHROMATOGRAPHY, vol. 18, no. 4, 13 February 1995, NEW YORK, pages 713-723, XP000646352 SHIMADA K. ET AL: "Studies on neurosteroids. I. retention behavior of derivatized 17-oxosteroids using high performance liquid chromatography" see page 715, paragraph 1; figure 1 ---	1,2,11
A	US 4 507 289 A (COLEMAN D.L. ET AL) 26 March 1985 cited in the application see column 3, line 14 - line 22 see column 3, line 55 - line 58 -----	3-5

# INTERNATIONAL SEARCH REPORT

...formation on patent family members

International Application No

PCT/US 96/18189

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 477107 A	25-03-92	FR 2667070 A WO 9204917 A	27-03-92 02-04-92
WO 9204888 A	02-04-92	US 5120720 A AU 8726891 A	09-06-92 15-04-92
EP 0149220 A	24-07-85	US 4666898 A AU 571626 B AU 3720184 A DE 3446886 A	19-05-87 21-04-88 04-07-85 18-07-85
EP 0261891 A	30-03-88	NONE	
US 4507289 A	26-03-85	DE 3474510 A EP 0148436 A	17-11-88 17-07-85